

Please review before being vaccinated

These individuals can't be vaccinated:

- Pfizer: Less than 12 years old- ***The vaccine is not approved for anyone less than 12 years old***
- Moderna: Less than 18 years old- ***The vaccine is not approved for anyone less than 18 years old***
- Moderate to severe illness on appointment date- ***Wait until feeling better.***
- COVID-19 positive and having symptoms- ***Wait until you are feeling better and out of isolation period.***
- Currently in a quarantine period due to exposure to someone with COVID- ***Come back when you are out of quarantine.***
- Received monoclonal antibodies or convalescent plasma as part of COVID-19 treatment in the past 90 days- ***Wait until 90 days after the treatment.***
- History of myocarditis or pericarditis prior to or after any dose of COVID-19 vaccine- ***Required to have approval of your medical provider prior to vaccination.***
- Received a COVID-19 vaccine not authorized or approved for use in the U.S.- ***Required to have approval of your medical provider prior to vaccination.***
- Allergic reaction to any brand of COVID-19 vaccine- ***should consult with an allergist for approval prior to vaccination***
- Allergic reaction to any component of the vaccine, including any of the following:
 - Polyethylene glycol (PEG), which is found in some medications, such as laxatives and preparations for colonoscopy procedures.
 - Polysorbate, which is found in some vaccines, film coated tablets, and intravenous steroids.
 - A vaccine or injectable therapy that contains multiple components, one of which is a COVID-19 vaccine component, but it is not known which component elicited the immediate reaction.

These individuals can be vaccinated, but are made aware of the risks/limitations:

- Less than 18 years old- ***Written Parent/guardian approval is required prior to vaccination.***
- Impaired immune responsiveness, whether due to use of immunosuppressive agents (irradiation, corticosteroids, etc.), HIV infection, or other causes- ***May have a reduced antibody response to active immunization.***
- Currently have dermal fillers- ***Swelling around the filler can occur; notify your healthcare provider if swelling develops at or near the site of the filler after vaccination.***
- Bleeding disorder or on blood thinner: ***There is an increased risk for bleeding at the injection site- If no intramuscular (IM) vaccine received since being diagnosed with bleeding disorder or taking a prescription blood thinner, check with your medical provider for approval to receive an IM vaccine.***
- If pregnant- ***There is no data on the safety of COVID-19 vaccine in pregnant women. There is a risk of adverse pregnancy outcome and studies are still ongoing. If you have questions or concerns about vaccination, discuss with your medical provider to make an informed decision.***
- If breastfeeding- ***There is no data on the safety of COVID-19 vaccine in breastfeeding women or on the effects on the breastfed infant or milk production/excretion. The vaccine does not contain a live virus, so no harm to the baby is expected. If questions or concerns about vaccination, discuss with your medical provider to make an informed decision.***
- History of multisystem inflammatory syndrome in children (MIS-C) or adults (MIS-A): ***There are no data on the safety and efficacy of COVID-19 vaccines in individuals with a history of multisystem inflammatory syndrome. The CDC recommends considering delaying vaccination until recovered from the illness and for 90 days after the date of diagnosis. You may want to discuss with your medical provider to assist with decisions about the use of a COVID-19 vaccine.***
- Received a first dose of another brand of vaccine and were unable to finish the series with the same brand of COVID-19 vaccine- ***All doses should be given with the same brand of vaccine.*** If unable to determine the previous product given, the available product can be given.